

The Consequences of Non-Normality

I. Hip^{a*}, Th. Lippert^a, H. Neff^b, K. Schilling^a and W. Schroers^a

^aDepartment of Physics, University of Wuppertal, D-42097 Wuppertal, Germany

^bNIC, c/o Research Center Jülich, D-52425 Jülich and DESY, D-22603 Hamburg, Germany

The non-normality of Wilson-type lattice Dirac operators has important consequences — the application of the usual concepts from the textbook (hermitian) quantum mechanics should be reconsidered. This includes an appropriate definition of observables and the refinement of computational tools. We show that the truncated singular value expansion is the optimal approximation to the inverse operator D^{-1} and we prove that due to the γ_5 -hermiticity it is equivalent to γ_5 times the truncated eigenmode expansion of the hermitian Wilson-Dirac operator.

1. INTRODUCTION

Wilson-type lattice Dirac operators are non-normal, i.e.

$$[D, D^\dagger] \neq 0, \quad (1)$$

and this fact has several important consequences. The essential difference to normal operators is that they are not diagonalizable by a unitary transformation. Instead they are diagonalizable (if at all) by a similarity transformation of the form

$$\text{diag}(\lambda_1, \dots, \lambda_N) = X^{-1} D X, \quad (2)$$

with $N = \dim(D)$. It follows that for a given eigenvalue λ_i there are distinct left $\langle L_i|$ and right $|R_i\rangle$ eigenvectors

$$D|R_i\rangle = \lambda_i|R_i\rangle, \quad (3)$$

$$\langle L_i|D = \lambda_i\langle L_i|, \quad (4)$$

in the sense that (in general) $\langle L_i| \neq (|R_i\rangle)^\dagger$.

This fact must be taken into account when performing the spectral decomposition

$$D = \sum_{i=1}^N \lambda_i |R_i\rangle \langle L_i| \neq \sum_{i=1}^N \lambda_i |R_i\rangle \langle R_i|, \quad (5)$$

but also raises the question of an appropriate generalization of the other concepts familiar from the

*Email: hip@theorie.physik.uni-wuppertal.de

textbook (hermitian) quantum mechanics. For example, if an observable in the continuum is defined as $\langle \psi_i|A|\psi_i\rangle$, where $|\psi_i\rangle$ is an eigenstate of the continuum operator D — what is the proper generalization for the case where the left and the right eigenvectors are distinct?

In [1] we advocate the use of $\langle L_i|A|R_i\rangle$ instead of $\langle R_i|A|R_i\rangle$ as a kind of improved observable. Note that in lattice field theory this question is not of importance in the continuum limit, because the difference between the two definitions disappears. However, this question was also raised in the domain of solid state physics — see [2] for an interesting discussion.

2. TRUNCATED EIGENMODE EXPANSION

In this contribution we want to examine a different problem, which is also connected to the non-normality of the Wilson-type lattice Dirac operators. The idea to approximate the inverse operator D^{-1} by just a few eigenmodes with eigenvalues close to zero is physically appealing. One hopes that the lowest modes dominate physics and that the truncated eigenmode expansion

$$D_k^{-1} = \sum_{i=1}^{k \ll N} \frac{1}{\lambda_i} |R_i\rangle \langle L_i| \quad (6)$$




Figure 1. Correlation function for several values of Δt (Δt increases as one steps down from the top curve) plotted versus the number of eigenmodes used to approximate the inverse of the Wilson-Dirac operator — for a given k , the correlation function is computed from D_k^{-1} , as defined in (6). The dimension of D is 512, so the final points give the values for the exact inverse of D^{-1} .

with the ordering

$$\frac{1}{|\lambda_1|} \geq \frac{1}{|\lambda_2|} \geq \dots \geq \frac{1}{|\lambda_k|} \geq \dots \geq \frac{1}{|\lambda_N|} \quad (7)$$

will be a fair approximation of the proper inverse $D^{-1} = D_N^{-1}$.

However, in [3] it was found that such an expansion is unstable and that it does not uniformly saturate by increasing k . This is illustrated in Fig. 1 where the values of the correlation function for different temporal distances Δt are plotted versus k . (The illustration is just for two dimensional QED on a 16^2 lattice for $\beta = 2$, but it should be stressed that for the discussion which follows this is irrelevant — the arguments are pure linear algebra and they will be valid for any physical system and parameter set.)

By a closer examination of pseudospectra (for more details about pseudospectra see [5]) of D , we identified that the spikes in Fig. 1 correspond essentially to the instabilities due to the non-normality of D . For non-normal operators the truncated eigenmode expansion in the original

biorthogonal basis seems to be problematic. However, one can do better than that.

3. SINGULAR VALUE DECOMPOSITION

For any matrix A there exists a so-called singular value decomposition (SVD)

$$A = U \Sigma V^\dagger = \sum_i \sigma_i |u_i\rangle \langle v_i|, \quad (8)$$

where U and V are unitary and Σ is the diagonal matrix with positive or zero entries on the diagonal — the “singular values” σ_i . We assume the following ordering:

$$\sigma_1 \geq \sigma_2 \geq \dots \sigma_N \geq 0. \quad (9)$$

Among other interesting properties, one can prove the following theorem (Th. 2.5.3. in [4]):

If $k < r = \text{rank}(A)$ and

$$A_k = \sum_{i=1}^k \sigma_i |u_i\rangle \langle v_i| \quad (10)$$

then

$$\min_{\text{rank}(B)=k} \|A - B\|_2 = \|A - A_k\|_2 = \sigma_{k+1}, \quad (11)$$

i.e., A_k is the closest matrix to A that has rank k . In other words, the truncation of SVD is the optimal approximation to the original matrix for a given k .

Actually, the drawback is that for a general matrix A one needs two sets of vectors: $|u\rangle$ -s and $|v\rangle$ -s. However, the situation is better for the Wilson-type lattice Dirac operators (and their inverses). Using the γ_5 -hermiticity property

$$D^\dagger = \gamma_5 D \gamma_5 \quad (12)$$

we are going to prove that the truncated SVD of D^{-1} is identical to

$$D_k^{-1} = \gamma_5 \sum_{i=1}^k \frac{1}{\mu_i} |w_i\rangle \langle w_i| = \gamma_5 H_k^{-1}, \quad (13)$$

with the ordering according to the absolute values of inverse μ -s:

$$\frac{1}{|\mu_1|} \geq \frac{1}{|\mu_2|} \geq \dots \geq \frac{1}{|\mu_k|} \geq \dots \frac{1}{|\mu_N|}. \quad (14)$$




Figure 2. Correlation function for several temporal distances Δt plotted versus the truncated singular value expansion, as defined by (13). This time, the number of eigenmodes refers to the modes of the hermitian Wilson-Dirac operator H used to construct H_k^{-1} .

The μ -s and $|w\rangle$ -s are the eigenvalues and eigenvectors of the hermitian Wilson-Dirac operator

$$H = \gamma_5 D = W M W^\dagger = \sum_{i=1}^N \mu_i |w_i\rangle \langle w_i|. \quad (15)$$

The hermiticity of H follows from the γ_5 -hermiticity property (12). Thus H is diagonalizable by a unitary transformation W and has real eigenvalues μ_i , ordered on the diagonal of the diagonal matrix M .

Per construction, $D^\dagger D$ is a hermitian matrix and it is easy to see that

$$D^\dagger D = W M W^\dagger W M W^\dagger = W M^2 W^\dagger. \quad (16)$$

However, by using the singular value decomposition (8) of D it follows that

$$D^\dagger D = V \Sigma U^\dagger U \Sigma V^\dagger = V \Sigma^2 V^\dagger. \quad (17)$$

As V is unitary and Σ^2 diagonal, one must have (with appropriate ordering)

$$\sigma_i^2 = \mu_i^2 \quad \Rightarrow \quad \sigma_i = |\mu_i|, \quad (18)$$

i.e., the singular values of D are just the absolute values of the eigenvalues of the hermitian Wilson-Dirac operator H . Also, the $|v\rangle$ -s should correspond to the $|w\rangle$ -s (there is an arbitrary phase

factor, but it cancels in (13)). Finally, we can rewrite:

$$D = U \Sigma V^\dagger = U \text{sign}(M) M W^\dagger = \gamma_5 W M W^\dagger \quad (19)$$

which makes it obvious that

$$U = \gamma_5 W \text{sign}(M) \Rightarrow |u_i\rangle = \text{sign}(\mu_i) \gamma_5 |w_i\rangle. \quad (20)$$

Hence, the truncated singular value expansion of D is

$$\begin{aligned} D_k &= \sum_{i=1}^k \sigma_i |u_i\rangle \langle v_i| = \sum_{i=1}^k \sigma_i \text{sign}(\mu_i) \gamma_5 |w_i\rangle \langle w_i| \\ &= \gamma_5 \sum_{i=1}^k \mu_i |w_i\rangle \langle w_i| = \gamma_5 H_k. \end{aligned} \quad (21)$$

As D^{-1} also satisfies the γ_5 -hermiticity property (12), the same reasoning applies and (13) follows.

4. CONCLUSION

Fig. 2 illustrates that, as expected, by using the truncated singular value expansion, non-normal artefacts disappear and the saturation as a function of k is superior to the expansion in the eigenmodes of D . This approach has already been successfully used in a realistic QCD setting [3].

Acknowledgments: I. H. thanks the DOE's Institute of Nuclear Theory at the University of Washington for its hospitality and the DOE for partial support during the completion of this work. W. S. is supported by the DFG Graduiertenkolleg "Feldtheoretische Methoden in der Elementarteilchentheorie und Statistischen Physik".

REFERENCES

1. I. Hip, T. Lippert, H. Neff, K. Schilling and W. Schroers, hep-lat/0105001.
2. N. Hatano and D. R. Nelson, Phys. Rev. B **58**, 8384 (1998), cond-mat/9805195.
3. H. Neff, N. Eicker, T. Lippert, J. W. Negele and K. Schilling, hep-lat/0106016.
4. G. H. Golub and C. F. van Loan, *Matrix Computations*, 3rd ed., John Hopkins University Press, Baltimore and London, 1996.
5. <http://web.comlab.ox.ac.uk/projects/pseudo-spectra/>